



## ***The Use and Application of Salt in Aquaculture***

LaDon Swann  
*Illinois-Indiana Sea Grant Program*  
Purdue University

Scott Fitzgerald, D.V.M.  
*Animal Disease Diagnostic Laboratory*  
Purdue University

### **Description**

Everyone is aware that salt is commonly used for flavoring and preserving foods. Did you know that salt is also one of most commonly used drugs in aquaculture? In fact, it is sometimes referred to as the aspirin of aquaculture.

Salt in its chemical form is sodium chloride (NaCl) and, as an approved chemical for food fish, requires no withdrawal time before marketing.

Many forms of salt are used, including table, meat-curing, pickling, and rock salt. Of these, the most commonly used and least expensive form is the meat-curing variety. When used properly, salt is employed in treating many external protozoans including *Costia*, *Epistylis*, *Trichodina*, *Chilodonella*, and the flukes *Dactylogyrus* and *Gyrodactylus*. Salt is used to treat bacterial gill infections and relieve stress during handling and transport. It also is effective for treating stress associated with nitrite poisoning in ponds.

Before any salt treatment is attempted a few general guidelines are suggested.

1. Use sensitive and accurate scales when calculating doses for treatment of small volumes of water contained in hauling or holding tanks. "Guesstimating" only ends in disaster.
2. Know the volumes of your tanks, pads, etc., beforehand. It is advisable to calculate these and have those values in a convenient location for ready use.
3. Perform a test treatment on a few fish before attempting large-scale treatment. Salt like other chemicals, reacts differently among different species and water qualities.

4. Prepare to remove fish or flush out salt baths with fresh water when fish first show signs of stress.

Treatment procedures involve calculating the volume of the water to treat, calculating the dosage of salt to apply, deciding which treatment method to use, and determining the rate of salt to apply for specific diseases. Each of these steps is outlined in the following sections.

### **Treatment Methods**

The method of salt application depends on the disease organism, the fish, and the size and type of the aquaculture unit. Treatment methods include short-term dips, prolonged baths, and indefinite treatments.

**Dip treatments** involve very strong solutions to which fish are exposed for short periods of time, usually 30 seconds to 1 minute.

**Prolonged baths** are useful for treating fish in small tanks that can be rapidly flushed. Strong solutions of salt are added to the water. Fish are held in this salt solution with aeration from 30-60 minutes or until they show signs of stress.

**Indefinite treatments** are used when transporting or handling fish, or when dealing with large volumes of water, such as ponds, where low concentrations of salt are used indefinitely.

### **Calculation of Volumes**

Knowing the volumes of your tanks and ponds beforehand is essential. Measurements used to determine volume are usually in feet and/or inches. The most common shapes of culture tank raceways are square, rectangular, or round tanks. Square or

rectangular pond volumes are calculated in the same manner as square or rectangular tanks except an average pond depth is used. The method used to calculate volume of each is given below.

**Volume of a Square or Rectangular Raceway, Tank, or Pond**

$$\text{Volume (Vol.)} = \text{Length} \times \text{Width} \times \text{Depth}$$

**Example 1:** A rectangular tank is 12 feet (ft.) long x 3 ft. wide and is 3 ft deep. What is its volume?

$$\begin{aligned} \text{Vol.} &= 12 \text{ ft.} \times 3 \text{ ft.} \times 3 \text{ ft.} \\ \text{Vol.} &= 108 \text{ cubic feet (ft.}^3) \end{aligned}$$

**Volume of a Round Tank**

$$\text{Vol.} = 3.14 \times (\text{radius} \times \text{radius}) \times \text{depth}$$

**Example 2:** A round tank is 12 ft. in diameter and 4.5 ft. deep. What is its volume? (Radius= 0.5 x diameter.)

$$\begin{aligned} \text{Vol.} &= 3.14 \times (\text{radius} \times \text{radius}) \times \text{depth} \\ \text{Vol.} &= 3.14 \times (6 \text{ ft.} \times 6 \text{ ft.}) \times 4.5 \text{ ft.} \\ \text{Vol.} &= 3.14 \times 36 \text{ ft.}^2 \times 4.5 \text{ ft.} \\ \text{Vol.} &= 508.7 \text{ ft.}^3 \end{aligned}$$

**Calculation of Dosages**

Once the volume is calculated in cubic feet, the gallons are determined using these conversions:

$$\begin{aligned} 1 \text{ ft.}^3 &= 7.48 \text{ gallons (gal.)} \\ 1 \text{ acre-foot (1 surface acre} &\times 1 \text{ ft. deep)} \\ &= 325,851 \text{ gal.} \\ 1 \text{ liter(1)} &= 0.26 \text{ gal.} \end{aligned}$$

Other useful conversions:

$$\begin{aligned} 1 \text{ pound (lb.)} &= 454 \text{ grams (g)} \\ 1,000 \text{ g} &= 1 \text{ kilogram (kg)} \end{aligned}$$

One of the most commonly used units of measure in aquiculture is the part-per-million, commonly referred to as ppm. If you calculate in percent, remember that 1% equals 10,000 ppm. The amounts of salt added to various volumes that result in 1 ppm concentrations are listed below.

$$\begin{aligned} 1 \text{ ppm equals:} \\ 2.7 \text{ lb./acre-ft.} \\ 0.0283 \text{ g/ft.}^3 \\ 0.00378 \text{ g/gal.} \\ 1.0 \text{ milligram(mg)/1} \end{aligned}$$

**Example 3:** How much salt is needed to make a 0.5% solution using an indefinite treatment in a 100-gal. transport tank?

$$\begin{aligned} 0.5\% &= 5,000 \text{ ppm} \\ \text{Salt needed} &= 0.00378 \text{ g/gal.} \times 100 \text{ gal.} \\ &\times 5000 \text{ ppm} \\ \text{Salt needed} &= 1,890 \text{ g or 4.2 lbs.} \end{aligned}$$

**Specific Treatment Rates**

Specific treatments using salt are given in Table 1.

<b>Table 1. Specific treatment rates and methods of using salt for treating various diseases or as a remedial treatment of stress.</b>	
<b>( To Control</b>	<b>Concentration and duration of treatment. )</b>
External parasites of brood fish:	30,000 ppm (3%) as a quick dip (15 seconds) before Stocking.
External parasites <i>Costia</i> , <i>Epistylis</i> , <i>Trichodina</i> , and <i>Chilodonella</i> and the flukes <i>Dactylogyrus</i> and <i>Gyrodactylus</i> :	10,000-30,000 ppm (1-3%) prolonged treatment (30 minutes or until fish show signs of stress) or 1,000-2,000 ppm in hauling tanks as an indefinite treatment
Nitrite poisoning or "Brown Blood Disease:"	50 pounds of salt per acre-foot when nitrite levels exceed 0.5 ppm.
Stress during transport and while handling:	Indefinite treatment using 1,000-10,000 ppm (0.1-1.0%).